

EQUITOP PRONUTRIN®

Equitop Pronutrin® maintains gastric health and provides extra nutrition in various conditions, e.g. loss of weight.

Equitop Pronutrin® contains 77% Apolectol®, a patented combination of fruit and vegetable fibre, glycerol and lecithin

When to use Equitop Pronutrin®?

Equitop Pronutrin® can be administered prior to, or during periods: when high performance is required, when feeding high levels of concentrates, changing types of feed, when feed is reduced, during transportation, weaning of foals or during rank conflicts in the herd. Equitop Pronutrin® can be given before or during competitions. It does not contain any doping relevant substances.

Feeding recommendations:

Standard feeding is 50g Equitop Pronutrin per 100kg bodyweight daily for a minimum of 28 days. (Scoop included measures from 50g to 125g in 25g increments). One container is sufficient for an average sized horse for 14 days. If you have questions regarding the suitability of use consult your veterinarian.

Racehorses and trotters:

The recommended quantity of Equitop Pronutrin® should be fed prior to training/racing. Other mature horses: The recommended quantity of Equitop Pronutrin® should be mixed with normal grain or supplement ration of the horse's diet and split equally between feedings. Example: If the horse is fed twice daily mix half the recommended daily quantity of Equitop Pronutrin® in the morning feed and the second half in the evening feed. If the horse is reluctant to eat Equitop Pronutrin® start with half the recommended daily quantity and gradually increase this over the next 2 to 3 days until recommended amount is reached. Foals: For suckling foals the recommended quantity of Equitop Pronutrin® is mixed with 3 times the quantity of lukewarm water. Leave the mixture for 2-3 minutes and then feed the foal by placing a syringe or bottle directly into its mouth.

Composition:

Dried fruit and vegetable pulp (66%), sugar (20%), apple extract (2%).

Additives:

Lecithin (9%), glycerol (2%), natural flavours (<1%), ethoxyquin (90 ppm).
Crude fibre 12%, crude fat 7.5%, crude protein 6%, crude ash 4%, magnesium 0.2%, sodium 0.0%.

Directions for Storage:

Always store with the lid properly closed. Product should be kept in a dry place and not exposed to direct sunlight. Caution: Keep out of reach of children. For animal use only. Not for human consumption

This is not a veterinary medicine which is subject to authorisation by the Irish Medicines Board.

Distributor in the UK/ROI:

Boehringer Ingelheim Vetmedica, Bracknell, Berks, RG12 8YS, UK
A division of Boehringer Ingelheim Ltd

Product responsible:

Boehringer Ingelheim Danmark A/S
Reg.no: 208-R854833